

## MOD3-MW300-Antepartum Complications

1. All of the following are risk factors associated with IUGR except:
  - a. Macrosomia
  - b. Maternal alcohol use
  - c. Renal disease
  - d. Maternal diabetes
  
2. The condition termed polyhydramnios is commonly associated with:
  - a. Elevated liver enzymes and hypertension
  - b. Maternal diabetes and multiple pregnancy
  - c. Trisomy 18 and IUGR
  - d. Fetal renal and hepatic malformation
  
3. Which of the following is TRUE about a Hydatidiform mole?
  - a. It is a fetal anomaly
  - b. It is characterized by a low hCG levels in the blood stream
  - c. It is the formation of a large mass of fluid filled cysts
  - d. It is a result of a genetic variant carried in the maternal DNA
  
4. The following are true about ectopic pregnancy except:
  - a. Ectopic pregnancy is a pregnancy that occurs whenever the blastocyst implants anywhere except in the endometrium of the uterine cavity.
  - b. Possible sites include: cervix, fallopian tubes, ovaries, and abdomen.
  - c. Early diagnosis is important for fetal survival
  - d. 95% are tubal.
  
5. Asymptomatic bacteriuria, cystitis and acute pyelonephritis are common during pregnancy. The following are true, except:
  - a. Urinary stasis contributes to the prevalence of UTIs in pregnancy
  - b. It is imperative that women not drink so much that their bladders are overextended in pregnancy.
  - c. Teaching hygiene, such as wiping from front to back is one way to help prevent UTI.
  - d. The most effective way to confirm an asymptomatic UTI is by a urine culture.
  
6. Which of the following is NOT an antepartum risk associated with multiple pregnancy?
  - a. Polyhydramnios
  - b. Prematurity
  - c. Twin-to-twin transfusion
  - d. Oligohydramnios

7. Which of the following is a possible result of hyperemesis gravidarum?
  - a. Electrolyte imbalance
  - b. Elevated blood pressure
  - c. Elevated liver enzymes
  - d. Glucosuria
  
8. Maternal Rubella infection during the first trimester is associated with which of the following?
  - a. Premature rupture of membranes
  - b. 50% chance of congenital defects of the fetus
  - c. Fetal rash and fever
  - d. Swollen neck glands of the fetus at birth
  
9. Iron deficient anemia is defined by which of the following values?
  - a. HGB 12 and HCT 36
  - b. HGB 11 and HCT 33
  - c. HGB 10 and HCT 30
  - d. HGB 12 and HCT 30
  
10. Which of the following symptoms is NOT associated with preeclampsia?
  - a. Proteinuria
  - b. Elevated blood pressure
  - c. Low platelets
  - d. Hematuria