

## MOD5-MW340- Postpartum Pharmacology

1. When is the most common time to administer RhoGAM postpartum?
  - a. Immediately after birth
  - b. Within 24 hours
  - c. Within 72 hours
  - d. Within 1-2 weeks
2. Which of the following drugs produces the effect of strong rhythmic contractions when administered IM postpartum?
  - a. Pitocin
  - b. Methergine
  - c. Lidocaine
  - d. Prostaglandins
3. Which of the following drugs should not be given to stop bleeding that occurs before the birth of the placenta?
  - a. Pitocin
  - b. Methergine
  - c. Misoprostol
  - d. IV fluids
4. Which of the following herbals actions is most desirable when treating a postpartum hemorrhage?
  - a. A uterine stimulant
  - b. A uterine relaxant
  - c. An anti-emetic
  - d. An anti-inflammatory
5. What is the most common dose of Methergine (Methylergonovine) that is given IM to help the uterus contract?
  - a. 0.2 mg
  - b. 2 mg
  - c. 10 IU
  - d. 20 IU
6. What is the most common dose of Pitocin (synthetic oxytocin) when given IM to help the uterus contract?
  - a. 0.2 mg
  - b. 2 mg
  - c. 10 IU
  - d. 20 IU
7. What anti-hemorrhage drug can be given orally, rectally or vaginally?
  - a. Pitocin
  - b. Methergine
  - c. Misoprostol

- d. Lidocaine
8. Which type of drugs is used most commonly in association with suturing?
- a. Analgesia
  - b. Local anesthesia
  - c. Anti-inflammatory
  - d. Hemostatic
9. Which type of drug is most commonly used to stimulate milk production?
- a. An oxytocic
  - b. A prolactin
  - c. A galactagogue
  - d. An emmenagogue
10. Which of the following remedies is an appropriate first treatment of sore nipples?
- a. Lanolin ointment
  - b. Over-the-counter anti-fungal cream
  - c. Gentian violet
  - d. Zinc oxide